<u> </u>				525 Rec'd PCT/PTO 20 NOV 2001			
			NT OF COMMERCE PATENT AND TRADE	ATTORNEY'S DOCKET NUMBER			
			R TO THE UNITED STA				
	D	DESIGNATED/ELECT	ΓED OFFICE (DO/EO/U	U S. APPLICATION NO. (IF KNOWN, SEE 37 CFR			
	CC	ONCERNING A FILI	NG UNDER 35 U.S.C. 3	09/700700			
TERN		NAL APPLICATION NO. CT/US99/10548	INTERNATIONAL FILING DA				
TLE C		ENTION	12 MAY 1999	21 MAY 1998			
		METHOD OF DIAGNOS 5) FOR DO/EO/US	SING, MONITORING, AND	D STAGING PROSTATE CANCER			
LI, S	hujat	th et al.					
nlica	nt here	exwith submits to the United S	totas Dagismatad/Elastad Office	(DO/FO/MON IL O.II)			
				(DO/EO/US) the following items and other information:			
. 2			f items concerning a filing under				
. [ncerning a filing under 35 U.S.C. 371.			
. ≥	☑ Thex	his is an express request to be xamination until the expiration	gin national examination proced n of the applicable time limit set	lures (35 U.S.C. 371(f)) at any time rather than delay in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).			
. 2				as made by the 19th month from the earliest claimed priority date			
. 🗵			plication as filed (35 U.S.C. 371				
	a.		th (required only if not transmitte				
	b.		by the International Bureau.	24.94			
	c.			ted States Receiving Office (RO/US).			
] A		al Application into English (35 U	- , , ,			
Þ		copy of the International Sea					
×		Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371 (c)(3))					
	a.		ith (required only if not transmitt				
	b.	ma.	by the International Bureau.	• · · · · · · · · · · · · · · · · · · ·			
	c.			ng such amendments has NOT expired.			
	d.						
] A1	translation of the amendmen	ts to the claims under PCT Article	le 19 (35 U.S.C. 371(c)(3)).			
×			eventor(s) (35 U.S.C. 371 (c)(4)).				
	_		liminary Examination Report (PC				
] A1			amination Report under PCT Article 36			
			nt(s) or information included:				
			tement under 37 CFR 1.97 and 1.				
<u></u>				in compliance with 37 CFR 3.28 and 3.31 is included.			
		FIRST preliminary amendm					
L.		SECOND or SUBSEQUEN	Γ preliminary amendment.				
		substitute specification.					
		change of power of attorney					
		ertificate of Mailing by Expre	ss Mail				
×	_	ther items or information:	and the second s				
	Ex	xecuted Verified Statement	Claiming Small Entity Status				
				Mail" Label No. #EL777534836US eposit - November 20, 2000			
			the United to Address indicated	certify that this paper is being deposited with d States Postal Service "Express Mail Post Office see" service under 37 CFR 1.10 on the date above and is addressed to the Assistant mer of Patents, Box PCT, Washington, D.C. 20231.			
	- 1		- Acc	- Angelowa			

		520 D	ec'd	DCT/DT 9/	NOVICE
U.S. APPLICATION NO. (IF KNOWN, SEE 37 CFR	INTERNATIONAL APPLICAT	TION NO.	it u	ATTORNEY	DOCKET NUMBER
-09/700700	PCT/US99/1054	18		DE	X-0105
21. The following fees are submitted:.				CALCULATION	S PTO USE ONLY
BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) - (Neither international preliminary examination					
international search fee (37 CFR 1.445(a)(2) rand International Search Report not prepared	paid to USPTO	\$1,00	00.00		
☐ International preliminary examination fee (37 USPTO but Internation Search Report prepare	CFR 1.482) not paid to d by the EPO or JPO	\$80	50.00		
International preliminary examination fee (37 but international search fee (37 CFR 1.445(a)((2)) paid to USPTO) \$7 1	10.00	:	
✓ International preliminary examination fee paid but all claims did not satisfy provisions of PC	Γ Article 33(1)-(4)	\$69	00.00		
☐ International preliminary examination fee paid and all claims satisfied provisions of PCT Arti	icle 33(1)-(4)		00.00		
	ATE BASIC FEE AMO			\$690.00	
Surcharge of \$130.00 for furnishing the oath or declar months from the earliest claimed priority date (37 CF)	ration later than \Box 20 FR 1.492 (e)).) 🗆 3	0	\$0.00	
CLAIMS NUMBER FILED	NUMBER EXTRA	RATE	3	**	
Total claims 11 - 20 =	0	x \$18.0	00	\$0.00	
Independent claims 5 - 3 =	2	x \$80.0	0	\$160.00	
Multiple Dependent Claims (check if applicable).	A DOVE CALCULAT	XONG.		\$270.00	******
Reduction of 1/2 for filing by small entity, if applical	ABOVE CALCULAT		=	\$1,120.00	
must also be filed (Note 37 CFR 1.9, 1.27, 1.28) (che	ck if applicable).	ement	×	\$560.00	
		<u> TOTAL</u>		\$560.00	
Processing fee of \$130.00 for furnishing the English t months from the earliest claimed priority date (37 CF)	ranslation later than 20 R 1.492 (f)).	31) +	\$0.00	
	TOTAL NATIONAL	FEE	=	\$560.00	
Fee for recording the enclosed assignment (37 CFR 1. accompanied by an appropriate cover sheet (37 CFR 3	21(h)). The assignment must b 3.28, 3.31) (check if applicable	e e).		\$0.00	
	TOTAL FEES ENCL	OSED	=	\$560.00	
				Amount to be: refunded	\$
			[charged	\$
☐ A check in the amount of	to cover the above fees is encl	losed.			
Please charge my Deposit Account No.	in the amount of			to cover the above	ve fees.
A duplicate copy of this sheet is enclosed.					
X Credit Card Payment for The Commissioner is hereby authorized to ch	arge any fees which may be req	uired, or cr	edit an	y overpayment	
to Deposit Account No. 501-619	A duplicate copy of this sheet is	enclosed.			
NOTE: Where an appropriate time limit under 37 1.137(a) or (b)) must be filed and granted to restore	CFR 1.494 or 1.495 has not be the application to pending st	een met, a tatus.	petitio	on to revive (37 CF	R
SEND ALL CORRESPONDENCE TO:					
LICATA, Jane Massey		SIGNATI	MCM 6	buch	
Licata & Tyrrrell P.C. 66 E. Main Street		SIGNATO			
Marlton, New Jersey 08053 US		Jane Ma	ssey I	Licata	
		NAME			
		32,257			
		REGISTR	ATIO	N NUMBER	
		20 Nove	mber	2000	
		DATE			
	1				

			Page 1 of 2
	EMENT (DECLARATION) CI FR 1.9(f) AND 1.27 (c)) - SMAL		
Serial No. Not Yet Assigned	Filing Date Herewith	Patent No.	Issue Date
Applicant/ ALI, Shuja Patentee:	th et al.		
Invention: A NOVEL	METHOD OF DIAGNOSING, MON	NITORING, AND STAGING PR	OSTATE CANCER
I hereby declare that I	am:		
	the small business concern identified		
🛚 an official of	the small business concern empowe	ered to act on behalf of the cond	ern identified below:
NAME OF CONCERN			
ADDRESS OF CONC	ERN: 3303 Octavius Drive, Santa Cla	ara, California 94043	
13 CFR 121.3-18, and of Title 35, United Sta not exceed 500 perso average over the previous basis during each of	the above-identified small business of reproduced in 37 CFR 1.9(d), for putes Code, in that the number of empins. For purposes of this statement, vious fiscal year of the concern of the pay periods of the fiscal year, one concern controls or has the power of both.	urposes of paying reduced fees ployees of the concern, includir (1) the number of employees on the persons employed on a full- and (2) concerns are affiliates	s under Section 41(a) and (b) ing those of its affiliates, does if the business concern is the time, part-time or temporary of each other when either,
I hereby declare that right identified above with r	rights under contract or law have bee egard to the above identified invention	en conveyed to and remain with on described in:	n the small business concern
★ the speciment ★ the speciment	ification filed herewith with title as list	ed above.	
☐ the applic	cation identified above.		
☐ the pater	nt identified above.		

If the rights held by the above-identified small business concern are not exclusive, each individual, concern or organization having rights to the invention is listed on the next page and no rights to the invention are held by any person, other than the inventor, who could not qualify as an independent inventor under 37 CFR 1.9(c) or by any concern which would not qualify as a small business concern under 37 CFR 1.9(d) or a nonprofit organization under 37 CFR 1.9(e).

				nave assigned, granted, convey, or license any rights in		
			organization ex or organization	kists. is listed below.		
FULL NAME _ADDRESS _						
FULL NAME _		Individual		Small Business Concern		Nonprofit Organization
FULL NAME		Individual		Small Business Concern		Nonprofit Organization
ADDRESS _		Individual		Small Business Concern		Nonprofit Organization
ADDRESS		Individual		Small Business Concern		Nonprofit Organization
invention aver I acknowledge entitlement to	ring to thei e the duty small ent	r status as s to file, in th ity status p	mall entities. (3 is application or rior to paying,	ach named person, concern 67 CFR 1.27) or patent, notification of any or at the time of paying, as a small entity is no longe	y change in s	status resulting in loss of of the issue fee or any
information ar willful false st Title 18 of the	nd belief ar atements a United Sta	e believed to and the like s ates Code, a	o be true; and so made are po nd that such wi	of my own knowledge are further that these statement unishable by fine or impriso illful false statements may jen is verified statement is direct	ts were made nment, or bo opardize the	e with the knowledge that th, under Section 1001 of
NAME OF PER			Mohan Iyer			
OTHER THAN			Vice Presiden	t, Business Development		
ADDRESS OF	PERSON S	SIGNING:	diaDexus, Inc	·		Add
			3303 Octaviu	s Drive		
			Santa Clara,	California 95054		
SIGNATURE:		Uoha	5. 9	₽ _V DATE	E: <u>///</u>	7/00

MOLECLES SO NOA SOOL

PCT/US99/10548

A NOVEL METHOD OF DIAGNOSING, MONITORING, AND STAGING PROSTATE CANCER

FIELD OF THE INVENTION

This invention relates, in part, to newly developed 5 assays for detecting, diagnosing, monitoring, staging, and prognosticating cancers, particularly prostate cancer.

BACKGROUND OF THE INVENTION

Cancer of the prostate is the most prevalent malignancy in adult males, excluding skin cancer, and is an 10 increasingly prevalent health problem in the United States. In 1996, it was estimated that in the United States, 41,400 deaths would result from this disease, indicating that prostate cancer is second only to lung cancer as the most common cause of death in the same population. If diagnosed 15 and treated early, when the cancer is still confined to the prostate, the chance of cure is significantly higher.

Treatment decisions for an individual are linked to the stage of prostate cancer present in that individual. A common classification of the spread of prostate cancer was 20 developed by the American Urological Association (AUA). AUA classification divides prostate tumors into four stages, Stage A, microscopic cancer within prostate, is A to D. further subdivided into stages A1 and A2. Sub-stage A1 is a well-differentiated cancer confined to one site within the Treatment is generally observation, radical 25 prostate. prostatectomy, or radiation. Sub-stage A2 is a moderately to poorly differentiated cancer at multiple sites within the Treatment is radical prostatectomy or radiation. Stage B, palpable lump within the prostate, is further 30 subdivided into stages B1 and B2. In sub-stage B1, the cancer forms a small nodule in one lobe of the prostate. stage B2, the cancer forms large or multiple nodules, or occurs in both lobes of the prostate. Treatment for both substages B1 and B2 is either radical prostatectomy or radiation.

- 2 -

Stage C is a large cancer mass involving most or all of the prostate and is further subdivided into two stages. In substage C1, the cancer forms a continuous mass that may have extended beyond the prostate. In sub-stage C2, the cancer forms a continuous mass that invades the surrounding tissue. Treatment for both these sub-stages is radiation with or without drugs. The fourth stage is metastatic cancer and is also subdivided into two stages. In sub-stage D1, the cancer appears in the lymph nodes of the pelvis. In sub-stage D2, the cancer involves tissues beyond lymph nodes. Treatment for both these sub-stages is systemic drugs to address the cancer as well as pain.

However, current prostate cancer staging methods are limited. As many as 50% of prostate cancers initially staged as A2, B, or C are actually stage D, metastatic: Discovery of metastasis is significant because patients with metastatic cancers have a poorer prognosis and require significantly different therapy than those with localized cancers. The five year survival rates for patients with localized and metastatic 20 prostate cancers are 93% and 29%, respectively.

Accordingly, there is a great need for increasingly sensitive methods for the staging of a cancer in a human to determine whether or not such cancer has metastasized and for monitoring the progress of a cancer in a human.

In the present invention, methods are provided for detecting, diagnosing, monitoring, staging and prognosticating cancers, particularly prostate cancer via seven (7) Prostate Specific Genes (PSG). The seven PSGs refer, among other things, to native proteins expressed by the genes comprising the polynucleotide sequences of any of SEQ ID NO: 1, 2, 3, 4, 5, 6 or 7. In the alternative, what is meant by the seven PSGs as used herein, means the native mRNAs encoded by the genes comprising any of the polynucleotide sequences of SEQ ID NO: 1, 2, 3, 4, 5, 6 or 7 or levels of the genes comprising

- 3 -

any of the polynucleotide sequences of SEQ ID NO: 1, 2, 3, 4, 5, 6 or 7.

Other objects, features, advantages and aspects of the present invention will become apparent to those of skill in the art from the following description. It should be understood, however, that the following description and the specific examples, while indicating preferred embodiments of the invention, are given by way of illustration only. Various changes and modifications within the spirit and scope of the disclosed invention will become readily apparent to those skilled in the art from reading the following description and from reading the other parts of the present disclosure.

SUMMARY OF THE INVENTION

Toward these ends, and others, it is an object of the present invention to provide a method for diagnosing the presence of prostate cancer in a patient which comprises measuring levels of PSG in a sample of cells, tissue or bodily fluid from the patient and comparing the measured levels of PSG with levels of PSG in preferably the same cells, tissue, or bodily fluid type of a control, wherein an increase in the measured PSG levels in the patient versus levels of PSG in the control is associated with prostate cancer.

Another object of the present invention is to provide a method of diagnosing metastatic prostate cancer in 25 a patient which comprises measuring PSG levels in a sample of cells, tissue, or bodily fluid from the patient and comparing the measured PSG levels with levels of PSG in preferably the same cells, tissue, or bodily fluid type of a control, wherein an increase in measured PSG levels in the patient versus 30 levels of PSG in the control is associated with a cancer which has metastasized.

Another object of the present invention is to provide a method of staging prostate cancer in a patient which comprises identifying a patient having prostate cancer,

measuring levels of PSG in a sample of cells, tissues, or bodily fluid obtained from the patient, and comparing the measured PSG levels with levels of PSG in preferably the same cells, tissue or bodily fluid type of a control. An increase 5 in measured PSG levels in the patient versus PSG levels in the control can be associated with a cancer which is progressing while a decrease or equivalent level of PSG measured in the patient versus the control can be associated with a cancer which is regressing or in remission.

Another object of the present invention is to provide a method of monitoring prostate cancer in a patient for the onset of metastasis. The method comprises identifying a patient having prostate cancer that is not known to have metastasized, periodically measuring levels of PSG in a sample 15 of cells, tissues, or bodily fluid obtained from the patient, and comparing the measured PSG levels with levels of PSG in preferably the same cells, tissue, or bodily fluid type of a control, wherein an increase in measured PSG levels versus control PSG levels is associated with a cancer which has 20 metastasized.

Yet another object of the present invention is to provide a method of monitoring the change in stage of prostate cancer in a patient which comprises identifying a patient having prostate cancer, periodically measuring levels of PSG 25 in a sample of cells, tissue, or bodily fluid obtained from the patient, and comparing the measured PSG levels with levels of PSG in preferably the same cells, tissues, or bodily fluid type of a control wherein an increase in measured PSG levels versus the control PSG levels is associated with a cancer 30 which is progressing and a decrease in the measured PSG levels versus the control PSG levels is associated with a cancer which is regressing or in remission.

Other objects, features, advantages and aspects of the present invention will become apparent to those of skill 35 in the art from the following description. It should be

- 5 -

understood, however, that the following description and the specific examples, while indicating preferred embodiments of the invention, are given by way of illustration only. Various changes and modifications within the spirit and scope of the disclosed invention will become readily apparent to those skilled in the art from reading the following description and from reading the other parts of the present disclosure.

DESCRIPTION OF THE INVENTION

The present invention relates to diagnostic assays 10 and methods, both quantitative and qualitative for detecting, diagnosing, monitoring, staging, and prognosticating cancers by comparing levels of PSG measured in a patient with levels of PSG in a control. What is meant by "levels of PSG" as used herein, means levels of the native protein expressed by the 15 gene comprising the polynucleotide sequence of any of SEQ ID NO: 1, 2, 3, 4, 5, 6 or 7. In the alternative, what is meant by "levels of PSG" as used herein, is levels of the native mRNA encoded by the gene comprising any of the polynucleotide sequence of SEQ ID NO: 1, 2, 3, 4, 5, 6 or 7 or levels of the 20 gene comprising any of the polynucleotide sequence of SEQ ID NO: 1, 2, 3, 4, 5, 6 or 7. Such levels are preferably measured in at least one of cells, tissues and/or bodily fluids, and includes determination of both normal and abnormal levels of PSGs. Thus, for instance, a diagnostic assay in 25 accordance with the invention for diagnosing overexpression of PSG protein compared to control bodily fluids, cells, or tissue samples may be used to diagnose the presence of cancers, including prostate cancer. Any of the seven PSGs may be measured alone in the methods of the invention, 30 together or in various combinations of the seven PSGs.

By "control" it is meant a human patient without cancer and/or non cancerous samples from the patient, also referred to herein as a normal human control; in the methods for diagnosing or monitoring for metastasis, control may also

include samples from a human patient that is determined by reliable methods to have prostate cancer which has not metastasized.

All the methods of the present invention may 5 optionally include measuring the levels of other cancer markers as well as PSG. Other cancer markers, in addition to PSG, useful in the present invention will depend on the cancer being tested and are known to those of skill in the art. For example, simultaneous testing for increases in PSA as well as 10 increases in PSG are also within the scope of the present invention and believed to provide a higher level of assurance that such cancer being tested is metastatic or the onset of metastasis has occurred.

Diagnostic Assays

The present invention provides methods for diagnosing the presence of prostate cancer by analyzing for changes in levels of PSG in cells, tissues or bodily fluids compared with levels of PSG in cells, tissues or bodily fluids of preferably the same type from a normal human control, wherein an increase 20 in levels of PSG in the patient versus the normal human control is associated with the presence of prostate cancer. Without limiting the instant invention, typically, for a quantitative diagnostic assay a positive result indicating the patient being tested has cancer is one in which cells, 25 tissues, or bodily fluid levels of the cancer marker, such as PSG, are at least two times higher, and most preferably are at least five times higher, than in preferably the same cells, tissues, or bodily fluid of a normal human control.

The present invention also provides a method of 30 diagnosing metastatic prostate cancer in a patient having prostate cancer which has not yet metastasized for the onset of metastasis. In the method of the present invention, a human cancer patient suspected of having prostate cancer which may have metastasized (but which was not previously known to 35 have metastasized) is identified. This is accomplished by a variety of means known to those of skill in the art. For example, in the case of prostate cancer, patients are typically diagnosed with prostate cancer following traditional detection methods.

In the present invention, determining the presence of PSG in cells, tissues, or bodily fluid, is particularly useful for discriminating between prostate cancer which has not metastasized and prostate cancer which has metastasized.

Existing techniques have difficulty discriminating 10 between prostate cancer which has metastasized and prostate cancer which has not metastasized and proper treatment selection is often dependent upon such knowledge.

In the present invention, the cancer marker levels measured in such cells, tissue, or bodily fluid are PSGs, and are compared with levels of PSG in preferably the same cells, tissue, or bodily fluid type of a normal human control. That is, if the cancer marker being observed is just PSG in serum, this level is preferably compared with the level of PSG in serum of a normal human patient. An increase in the PSG in the patient versus the normal human control is associated with prostate cancer which has metastasized.

Without limiting the instant invention, typically, for a quantitative diagnostic assay a positive result indicating the cancer in the patient being tested or monitored 25 has metastasized is one in which cells, tissues, or bodily fluid levels of the cancer marker, such as PSG, are at least two times higher, and most preferable are at least five times higher, than in preferably the same cells, tissues, or bodily fluid of a normal patient.

30 Staging

The invention also provides a method of staging prostate cancer in a human patient.

The method comprises identifying a human patient having such cancer and analyzing a sample of cells, tissues, or bodily fluid from such patient for PSG. Then, the method

- 8 -

compares PSG levels in such cells, tissues, or bodily fluid with levels of PSG in preferably the same cells, tissues, or bodily fluid type of a normal human control sample, wherein an increase in PSG levels in the patient versus the normal human control is associated with a cancer which is progressing and a decrease in the levels of PSG is associated with a cancer which is regressing or in remission.

Monitoring

Further provided is a method of monitoring prostate 10 cancer in a human having such cancer for the onset of metastasis. The method comprises identifying a human patient having such cancer that is not known to have metastasized; periodically analyzing a sample of cells, tissues, or bodily fluid from such patient for PSG; and comparing the PSG levels in such cells, tissue, or bodily fluid with levels of PSG in preferably the same cells, tissues, or bodily fluid type of a normal human control sample, wherein an increase in PSG levels in the patient versus the normal human control is associated with a cancer which has metastasized.

Further provided by this invention is a method of monitoring the change in stage of prostate cancer in a human having such cancer. The method comprises identifying a human patient having such cancer; periodically analyzing a sample of cells, tissue, or bodily fluid from such patient for PSG; comparing the PSG levels in such cells, tissue, or bodily fluid with levels of PSG in preferably the same patient.

Monitoring such patient for onset of metastasis is periodic and preferably done on a quarterly basis. However, this may be more or less frequent depending on the cancer, the particular patient, and the stage of the cancer.

Assay Techniques

Assay techniques that can be used to determine levels of gene expression, such as PSG of the present invention, in a sample derived from a host are well-known to those of skill in the art. Such assay methods include radioimmunoassays,

curve.

WO 99/60162 PCT/US99/10548

- 9 -

reverse transcriptase PCR (RT-PCR) immunohistochemistry assays, in situ hybridization assays, competitive-binding assays, Western Blot analyses and ELISA assays. Among these, ELISAs are frequently preferred to 5 diagnose a gene's expressed protein in biological fluids. An ELISA assay initially comprises preparing an antibody, if not readily available from a commercial source, specific to PSG, preferably a monoclonal antibody. In addition a reporter antibody generally is prepared which binds specifically to 10 PSG. The reporter antibody is attached to a detectable reagent such as radioactive, fluorescent or enzymatic reagent, for example horseradish peroxidase enzyme or alkaline phosphatase.

To carry out the ELISA, antibody specific to PSG is 15 incubated on a solid support, e.g., a polystyrene dish, that binds the antibody. Any free protein binding sites on the dish are then covered by incubating with a non-specific protein such as bovine serum albumin. Next, the sample to be analyzed is incubated in the dish, during which time PSG binds 20 to the specific antibody attached to the polystyrene dish. Unbound sample is washed out with buffer. A reporter antibody specifically directed to PSG and linked to horseradish peroxidase is placed in the dish resulting in binding of the reporter antibody to any monoclonal antibody bound to PSG. 25 Unattached reporter antibody is then washed out. Reagents for peroxidase activity, including a colorimetric substrate are then added to the dish. Immobilized peroxidase, linked to PSG antibodies, produces a colored reaction product. The amount of color developed in a given time period is proportional to 30 the amount of PSG protein present in the sample. Quantitative

A competition assay may be employed wherein antibodies specific to PSG attached to a solid support and labeled PSG and a sample derived from the host are passed over

results typically are obtained by reference to a standard

- 10 -

the solid support and the amount of label detected attached to the solid support can be correlated to a quantity of PSG in the sample.

Nucleic acid methods may be used to detect PSG mRNA5 as a marker for prostate cancer. Polymerase chain reaction (PCR) and other nucleic acid methods, such as ligase chain reaction (LCR) and nucleic acid sequence based amplification (NASABA), can be used to detect malignant cells for diagnosis and monitoring of various malignancies. For example, reverse-10 transcriptase PCR (RT-PCR) is a powerful technique which can be used to detect the presence of a specific mRNA population in a complex mixture of thousands of other mRNA species. RT-PCR, an mRNA species is first reverse transcribed to complementary DNA (cDNA) with use of the enzyme reverse 15 transcriptase; the cDNA is then amplified as in a standard PCR RT-PCR can thus reveal by amplification the presence of a single species of mRNA. Accordingly, if the mRNA is highly specific for the cell that produces it, RT-PCR can be used to identify the presence of a specific type of 20 cell.

Hybridization to clones or oligonucleotides arrayed on a solid support (i.e., gridding) can be used to both detect the expression of and quantitate the level of expression of that gene. In this approach, a cDNA encoding the PSG gene is fixed to a substrate. The substrate may be of any suitable type including but not limited to glass, nitrocellulose, nylon or plastic. At least a portion of the DNA encoding the PSG gene is attached to the substrate and then incubated with the analyte, which may be RNA or a complementary DNA (cDNA) copy of the RNA, isolated from the tissue of interest.

Hybridization between the substrate bound DNA and the analyte can be detected and quantitated by several means including but not limited to radioactive labeling or fluorescence labeling of the analyte or a secondary molecule designed to detect the hybrid. Quantitation of the level of

gene expression can be done by comparison of the intensity of the signal from the analyte compared with that determined from known standards. The standards can be obtained by *in vitro* transcription of the target gene, quantitating the yield, and then using that material to generate a standard curve.

The above tests can be carried out on samples derived from a variety of patients' cells, bodily fluids and/or tissue extracts (homogenates or solubilized tissue) such as from tissue biopsy and autopsy material. Bodily fluids useful in the present invention include blood, urine, saliva, or any other bodily secretion or derivative thereof. Blood can include whole blood, plasma, serum, or any derivative of blood.

EXAMPLES

The present invention is further described by the following examples. These examples are provided solely to illustrate the invention by reference to specific embodiments. These exemplifications, while illustrating certain specific aspects of the invention, do not portray the limitations or circumscribe the scope of the disclosed invention.

EXAMPLE 1: PSGs

Searches were carried out and PSGs identified using the following Search Tools as part of the LIFESEQ® database available from Incyte Pharmaceuticals, Palo Alto, CA:

- 1. Library Comparison (compares one library to one other library) allows the identification of clones expressed in tumor and absent or expressed at a lower level in normal tissue.
- 2. Subsetting is similar to library comparison but 30 allows the identification of clones expressed in a pool of libraries and absent or expressed at a lower level in a second pool of libraries.

- 3. Transcript Imaging lists all of the clones in a single library or a pool of libraries based on abundance. Individual clones can then be examined using Electronic Northerns to determine the tissue sources of their component 5 ESTs.
- 4. Protein Function: Incyte has identified subsets of ESTs with a potential protein function based on homologies to known proteins. Some examples in this database include Transcription Factors and Proteases. Some leads were identified by searching in this database for clones whose component ESTs showed disease specificity.

Electronic subtractions, transcript imaging and protein function searches were used to identify clones, whose component ESTs were exclusively or more frequently found in libraries from specific tumors. Individual candidate clones were examined in detail by checking where each EST originated.

Table 1:

	SEQ ID	Clone ID #	Gene ID	
20	NO:		#	
	1	1550426	244673	Protein Function
				(Transcription Factors)
	2	1255804	14878	Subsetting
	3	1808432	255819	Subsetting
	4	3930803	none	Subsetting
25	5	645804	235032	Subsetting
	6	1862352	221558	Subsetting
	7	1450626	236019	Subsetting

EXAMPLE 2: Measurement of SEQ ID NO:1; Clone ID # 1550426;

Gene ID #244673 (pro101)

The example is carried out using standard techniques, which are well known and routine to those of skill in the art,

except where otherwise described in detail. Routine molecular biology techniques of the following example are carried out as described in standard laboratory manuals, such as Sambrook et al., MOLECULAR CLONING: A LABORATORY MANUAL, 2nd Ed.; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989).

Relative Quantitation of Gene Expression

Real-time quantitative PCR with fluorescent Taqman probes is a quantitative detection system utilizing the 5'10 3' nuclease activity of Taq DNA polymerase. The method uses an internal fluorescent oligonucleotide probe (Taqman) labeled with a 5' reporter dye and a downstream, 3' quencher dye. During PCR, the 5'-3' nuclease activity of Taq DNA polymerase releases the reporter, whose fluorescence can then be detected by the laser detector of the Model 7700 Sequence Detection System (PE Applied Biosystems, Foster City, CA, USA).

Amplification of an endogenous control is used to standardize the amount of sample RNA added to the reaction and normalize for Reverse Transcriptase (RT) efficiency. Either 20 cyclophilin, glyceraldehyde-3-phosphate dehydrogenase (GAPDH) or 18S ribosomal RNA (rRNA) is used as this endogenous control. To calculate relative quantitation between all the samples studied, the target RNA levels for one sample are used as the basis for comparative results (calibrator). 25 Quantitation relative to the "calibrator" is obtained using the standard curve method or the comparative method (User Bulletin #2: ABI PRISM 7700 Sequence Detection System).

To evaluate the tissue distribution, and the level of prol01 (SEQ ID NO:1) in normal and tumor tissue, total RNA was extracted from tumor and matched normal adjacent tissues and from unmatched tumor and normal tissues. Subsequently, first strand cDNA was prepared with reverse transcriptase and the polymerase chain reaction carried out using primers and Taqman probe specific to prol01 (SEQ ID NO:1). The results

were obtained using the ABI PRISM 7700 Sequence Detector. The absolute numbers are relative levels of expression of prol01 (SEQ ID NO:1) compared to the calibrator.

The absolute numbers are depicted in the following 5 Table 2 as relative levels of expression in 12 normal tissues of prol01 (SEQ ID NO:1) compared to kidney (calibrator). These RNA samples were generated by pooling samples from a particular tissue from different individuals.

Table 2: Relative levels of pro101 Expression in Pooled Samples

Tissue	NORMAL
Brain	1.2
Heart	2
Kidney	1
Liver	7.2
Lung	48.2
Mammary	2.5
Prostate	1418.4
Spleen	1.6
Small	1.9
Testis	57.3
Thymus	1.3
Uterus	7.6

The relative levels of expression in Table 2 show that for the 25 PSG pro101 (SEQ ID NO:1) mRNA expression is more than 20 fold higher in the pool of normal prostate compared with the other 11 normal tissue pools analyzed. These results demonstrate that mRNA expression of the PSG is highly specific for prostate.

The tissues shown in Table 2 correspond to pools of samples from different individuals. The tissues shown in the following Table 3 were obtained from individuals and are not pooled. Hence the values for mRNA expression levels shown in Table 2 cannot be directly compared to the values shown in Table 3.

15

10

20

The absolute numbers in Table 3 are relative levels of expression of pro101 (SEQ ID NO:1) compared to kidney (calibrator), in 60 pairs of matching samples. Each matching pair contains the cancer sample for a particular tissue and 5 the normal adjacent sample for that same tissue from the same individual. The results from 3 unmatched ovary tumor, 3 unmatched normal ovary, 1 unmatched mammary tumor and 1 unmatched normal mammary gland are also shown.

Table 3: Relative Levels of pro101 Expression in Individual Samples

	TISSUE	CANCER	MATCHING	UNMATCHED
	Prostate 1	103.9	0	
	Prostate 2	2219	84.2	
	Prostate 3	5048.2	3623.6	
15	Prostate 4	11052.3	2029.4	
	Prostate 5	229.1	41.1	
	Prostate 6	57.9	25.3	
	Prostate 7	58.5	57.069	
	Prostate 8	1074.6	610.8	
20	Prostate 9	32.7	79.3	
	Prostate 10	15.8	2.09	
	Prostate 11	436.4	438	
	Prostate 12	49.5	59.3	
	Prostate 13	128	56	
25	Bladder 1	0	0	
	Bladder 2	0	0	
	Bladder 3	0.7	0	
	Colon 1	0	0	
	Colon 2	0	0	
30	Colon 3	0	0	
	Colon 4	3.3	1.9	
	Colon 5	0.1	0.8	
	Colon 6	0	0	
	Lung 1	0	0	
35	Lung 2	0.5	1.6	
	Lung 3	1.4	2.1	
	Lung 4	0	0	
	Lung 5	0	0	
	Kidney 1	0	0	
40	Kidney 2	0	_0	
	Kidney 3	0	0	
	Kidney 4	0	0	
	Liver 1	1.5	5.7	
1 E	Liver 2	26.9	7.9	
45	Liver 3	0	0	

-		
 - 1	n	-

	Pancreas 1	0.9	0.9	
	Pancreas 2	3	Ô	
	Pancreas 3	0	0	
	Pancreas 4	0	0	
5	Pancreas 5	0	0	
	Stomach 1	0	0	
	Stomach 2	0	0	
	Stomach 3	0	0	
	Stomach 4	0	0	
10	Stomach 5	0	0	
	Sm Int 1	0	0	
	Sm Int 2	0	0	
	Testis 1	0	0	
	Mammary 1	4	0	
15	Mammary 2	5.6	0	
	Mammary 3	0.5	0	
	Mammary 4	0.4	0	
	Mammary 5	0.5		
	Mammary 6			0
20	Endo 1	1.6	7.6	
	Endo 2	0	0	
	Endo 3	0	0	
	Endo 4	0.3	0.2	
	Endo 5	5.8	5	
25	Uterus 1	0	0	
	Uterus 2	0	0	
	Uterus 3	0	0	
	Uterus 4	2.2	2.6	
	Ovary 1	1.4		
30	Ovary 2			11.6
	Ovary 3	1.5		
	Ovary 4		, <u></u>	22.9
	Ovary 5	0		1 0
	Ovary 6	<u> </u>		1.8

Among 128 samples in Table 3 representing 14 different tissues, the higher levels of expression are consistently in prostate tissues. These results confirm the tissue specificity results obtained with normal samples shown in Table 2. Table 2 and Table 3 represent a combined total of 140 samples in 18 human tissue types. Sixty-eight samples representing 13 different tissue types excluding prostate had no detected pro101 mRNA (Table 3). In 4 tissues (stomach small intestine kidney and testis) no pro101 (SEQ ID NO:1) mRNA was detected for any sample tested from individuals (Table 3). Expression of this PSG was detected in testis in the pooled normal sample (Table 3). The median expression in

25 prostate tissues.

WO 99/60162 PCT/US99/10548

- 17 -

prostate cancer samples in Table 3 is 166.5 units. Excluding Ovary 4 (Normal), only 1 sample in Table 3, Liver 2 (Cancer), is greater than 10% of this value.

Comparisons of the level of mRNA expression in 5 prostate tumor samples and the normal adjacent tissue from the same individuals are also shown in Table 3. The PSG pro101 (SEQ ID NO:1) is expressed at higher levels in 9 of 13 (69%) prostate cancer tissues (Prostate 1, 2, 3, 4, 5, 6, 8, 10 and 13) compared with the corresponding normal adjacent tissue. 10 The level of expression of this PSG is lower in prostate tumor compared to normal adjacent tissue in two samples (Prostate 9 and 12). Equivalent levels of expression were detected in two matched samples (Prostate 7 and 11). Previous mRNA expression analysis for genes coding for the diagnostic 15 markers PSA and PLA2 showed higher expression of the mRNA in 40% to 80% of the tumor samples compared to matching normal adjacent tissue. Higher expression in the tumor sample compared to the corresponding normal adjacent tissue is observed for Bladder 3, Colon 4, Liver 2, Pancreas 2, 20 Endometrium 5 and. Mammary 1, 2 and 3. Higher expression in the normal adjacent samples is observed for Colon 5, Lung 2, Lung 3, Liver 1, Endometrium 1 and Uterus 4. However, the levels detected are in most cases comparable amongst the different tissues and low compared to levels found in most

The high level of tissue specificity, plus the mRNA overexpression in 9 of 13 of the prostate tumor samples tested compared to the normal adjacent tissues are believed to make the PSG, pro101 (SEQ ID NO:1) a good diagnostic marker for detection of prostate cancer using mRNA.

- 18 -

WO 99/60162 PCT/US99/10548

What is Claimed is:

1. A method for diagnosing the presence of prostate cancer in a patient comprising:

- (a) measuring levels of PSG in a sample of cells, 5 tissue or bodily fluid obtained from the patient; and
- (b) comparing the measured levels of PSG with levels of PSG in a sample of cells, tissue or bodily fluid obtained from a control, wherein an increase in measured levels of PSG in the patient versus the PSG levels in the control is 10 associated with the presence of prostate cancer.
 - A method of diagnosing metastatic prostate 2. cancer in a patient comprising:
- (a) measuring levels of PSG in a sample of cells, 15 tissue, or bodily fluid obtained from the patient; and
- (b) comparing the measured levels of PSG with levels of PSG in a sample of cells, tissue, or bodily fluid obtained from a control, wherein an increase in measured PSG levels in the patient versus the PSG levels in the control is associated 20 with a cancer which has metastasized.
 - A method of staging prostate cancer in a patient comprising:
 - (a) identifying a patient suffering from prostate cancer;
- (b) measuring levels of PSG in a sample of cells, 25 tissue, or bodily fluid obtained from the patient; and
- (c) comparing the measured levels of PSG with levels of PSG in a sample of cells, tissue, or bodily fluid obtained from a control, wherein an increase in the measured levels of 30 PSG versus the levels of PSG in the control is associated with a cancer which is progressing and a decrease in the measured levels of PSG versus the levels of PSG in the control is associated with a cancer which is regressing or in remission.

WO 99/60162 PCT/US99/10548

- 19 -

4. A method of monitoring prostate cancer in a patient for the onset of metastasis comprising:

- (a) identifying a patient having prostate cancer that is not known to have metastasized;
- (b) periodically measuring PSG levels in samples of cells, tissue, or bodily fluid obtained from the patient; and
- (c) comparing the periodically measured levels of PSG with levels of PSG in cells, tissue, or bodily fluid obtained from a control, wherein an increase in any one of the periodically measured levels of PSG in the patient versus the levels of PSG in the control is associated with a cancer which has metastasized.
 - 5. A method of monitoring changes in a stage of prostate cancer in a patient comprising:
 - (a) identifying a patient having prostate cancer;
 - (b) periodically measuring levels of PSG in samples of cells, tissue, or bodily fluid obtained from the patient; and
- (c) comparing the measured levels of PSG with levels of PSG in a sample of the same cells, tissue, or bodily fluid of a control, wherein an increase in any one of the periodically measured levels of PSG versus levels of PSG in the control is associated with a cancer which is progressing in stage and a decrease in any one of the periodically measured levels of PSG versus the levels of PSG in the control is associated with a cancer which is regressing in stage or in remission.
 - 6. The method of claim 1, 2, 3, 4 or 5 wherein the PSG comprises SEQ ID NO:1.

Docket No. **DEX-0105**

Declaration and Power of Attorney For Patent Application English Language Declaration

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

A NOVEL METHOD OF DIAGNOSING, MONITORING, AND STAGING PROSTATE CANCER

the specification of whi	ch		
(check one)			
 □ is attached hereto. ☑ was filed on 12 Ma Application Number and was amended 	PCT/US99/10548	as United States Application No.	or PCT International
		(if applicable)	
I hereby state that I had including the claims, as	ave reviewed and u s amended by any a	nderstand the contents of the above in amendment referred to above.	dentified specification,
I acknowledge the dut known to me to be n Section 1.56.	ty to disclose to the material to patental	United States Patent and Trademark bility as defined in Title 37, Code of	Office all information Federal Regulations,
Section 365(b) of any any PCT International listed below and have	r foreign application application which d also identified belo r PCT International	under Title 35, United States Code, (s) for patent or inventor's certificate esignated at least one country other tw, by checking the box, any foreign a application having a filing date before	, or Section 365(a) of han the United States, pplication for patent or
Prior Foreign Applicati	ion(s)		Priority Not Claimed
(Number)	(Country)	(Day/Month/Year Filed)	
(Number)	(Country)	(Day/Month/Year Filed)	J
(Number)	(Country)	(Day/Month/Year Filed)	
PTO-SB-01 (9-95) (Modified)	www.	P02/REV02 Patent and Trademark	Office-U.S. DEPARTMENT OF COMMER

60/086,265	21 May 1998	
(Application Serial No.)	(Filing Date)	
(Application Serial No.)	(Filing Date)	
(Application Serial No.)	(Filing Date)	
hereby claim the henetit under :	76 II S 1 S AOTION 1711 OT	
Section 365(c) of any PCT Internance insofar as the subject matter of elunited States or PCT International U.S.C. Section 112, I acknowledgo Office all information known to metal information in the section of the sec	tional application designating ach of the claims of this application in the manner plus the duty to disclose to the late to be material to patentable between the filing date of	any United States application(s), on the United States, listed below and olication is not disclosed in the prior rovided by the first paragraph of 35 United States Patent and Trademark lity as defined in Title 37, C. F. R. the prior application and the national
Section 365(c) of any PCT Internance insofar as the subject matter of e United States or PCT International J.S.C. Section 112, I acknowledgo Office all information known to make the Section 1.56 which became available.	tional application designating ach of the claims of this application in the manner plus the duty to disclose to the late to be material to patentable between the filing date of	the United States, listed below and blication is not disclosed in the prior rovided by the first paragraph of 35 United States Patent and Trademark ility as defined in Title 37, C. F. R.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

(Filing Date)

(Application Serial No.)

(patented, pending, abandoned)

(Status)

(patented, pending, abandoned)

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith. (list name and registration number)



26259

PATENT TRADEMARK OFFICE

Send	Correspondence	to:
------	----------------	-----

Direct Telephone Calls to: (name and telephone number)
Jane Massey Licata or Kathleen A. Tyrrell - (856) 810-1515

Full name of sole or first inventor

Shujath Ali

Sole or first inventor's signature

Residence)

Santa Clara, California

Citizenship

US

Post Office Address

3475 Granada Avenue, Apt. 357

Santa Clara, California

Full name of second inventor, if any

Susana Salceda

Secon nvgntor's signature

Date

11/06/

2000

11-06-2000

Residence

San Jose, California

Citizenship

US

Post Office Address

4118 Cresendo Avenue

San Jose, California

Full name of third inventor, if any Yongming Sun	
Third, inventor's signature	Date
GRUS Su	11-6-2000
Residence San Jose, California	
Citizenship	
US Post Office Address	
869 S. Winchester Boulevard, Apt. 260	
San Jose, California	
Full name of fourth inventor, if any	
Robert Cafferkey	
Fourth inventor's signature 4	11-7-00 Date
Residence Mountain View, California	
Citizenship US	
Post Office Address	
651 Franklin Street, Apt. 4305	
Mountain View, California	
Full name of fifth inventor, if any	
Fifth inventor's signature	Date
Residence	
Citizenship	
Post Office Address	
Full name of sixth inventor, if any	
Sixth inventor's signature	Date
Residence	
Citizenship	
Post Office Address	

SEQUENCE LISTING

```
<110> Ali, Shujath
             Salceda, Susana
             Sun, Yangming
             Cafferkey, Robert
       <120> A Novel Method of Diagnosing, Monitoring and Staging
              Prostate Cancer
       <130> DEX-0034
       <140>
       <141>
       <150> 60/086,265
       <151> 1998-05-21
<160> 7
       <170> PatentIn Ver. 2.0
       <210> 1
       <211> 1936
       <212> DNA
       <213> Homo sapiens
       <220>
       <221> unsure
       <222> (1908)
       <400> 1
       aatggtatgc caacttaagt atttacaggg tggcccaaat agaacaagat gcactcgctg 60
       tgattttaag acaagetgta taaacagaac tccactgcaa gagggngggc cgggccagga 120
       gaateteege ttgteeaaga caggggeeta aggagggtet ceacactget getagggget 180
       gttgcatttt tttattagta gaaagtggaa aggcctcttc tcaacttttt tcccttgggc 240
       tggagaattt agaatcagaa gtttcctgga gttttcaggc tatcatatat actgtatcct 300
       gaaaggcaac ataattette etteeeteet titaaaatti tqtqtteett titqeagcaa 360
       ttactcacta aagggcttca ttttagtcca gatttttagt ctggctgcac ctaacttatg 420
        cctcgcttat ttagcccgag atctggtctt ttttntgtnt tttttttntt tccgtctccc 480
       caaagcttta tetgtettga etttttaaaa aagtttgggg geagattetg aattgggeta 540
        aaagacatgc atttttaaaa ctaggcaact tcttatttct ttcctttaaa aatacatagc 600
        attaaatccc aaatcctatt taaagacctg acagcttgag aaggtcacta ctgcatttat 660
        aggacettet ggtggttetg etgttaegtt tgaagtetga caateettga gaatetttge 720
       atgcagagga ggtaagaggt attggatttt cacagaggaa gaacacagcg cagaatgaag 780
        ggccaggctt actgaggctg tccagtggag ggctcatggg tgggacatgg aaaagaaggc 840
        agcctaggcc ctggggagcc cagtccactg agcaagcaag ggactgagtg agccttttgc 900
        aggaaaaggc taagaaaaag gaaaaccatt ctaaaacaca acaagaaact gtccaaatgc 960
```

ĭ

```
tttqqqaact qtqtttattq cctataatqq qtccccaaaa tqqqtaacct agacttcaqa 1020
gagaatgagc agagagcaaa ggagaaatct ggctgtcctt ccattttcat tctgttatct 1080
caggtgagct ggtagagggg agacattaga aaaaaatgaa acaacaaaac aattactaat 1140
gaggtacgct gaggcctggg agtctcttga ctccactact taattccgtt tagtgagaaa 1200
cettteaatt ttetttatt agaaggeea gettaetgtt ggtggeaaaa ttgeeaacat 1260
aagttaatag aaagttggcc aatttcaccc cattttctgt ggtttgggct ccacattgca 1320
atgttcaatg ccacqtqctq ctqacaccqa ccqqaqtact agccagcaca aaagqcaqqq 1380
tagoctgaat tgctttctgc tctttacatt tcttttaaaa taagcattta gtgctcagtc 1440
cetactgagt actettete teccetecte tgaatttaat tettteaaet tgeaatttge 1500
aaggattaca catttcactg tgatgtatat tgtgttgcag ngaaaagaaa aaagtgtctt 1560
tgtttaaaat tacttggttt qtgaatccat cttgcttttt ccccattgga actagtcatt 1620
aacccatctc tqaactggta qaaaaacatc tqaaqagcta gtctatcagc atctgacagg 1680
tgaattggat ggttctcaga accatttcac ccagacagcc tgtttctatc ctgtttaata 1740
aattagtttg ggttctctac atgcataaca aaccctgctc caatctgtca cataaaagtc 1800
tgtgacttga agtttagtca qcaccccac caaactttat ttttctatgt gttttttgca 1860
acatatgagt gttttgaaaa taaagtaccc atgtctttat taaaaaaanaa aaaaaagggc 1920
ggccgccgac tagtga
                                                                  1936
<210> 2
<211> 637
<212> DNA
<213> Homo sapiens
<400> 2
gtaggggcag acttactqcc ttqaacqaaa gacqatqgtc ctcgctcagc ctcactccaa 60
ttatgtteet ctaggtgggg caggtagggg qtccagette ctgettgetg gtggttcagg 120
teatgegtee ageettgtee ettetgaeet gggeeetaee eaeggggaaa tgtteeeata 180
gcagaagaat cagccccaca gtgcaggggt gtgttagtgg ggaacgggct ctgggctcct 240
gtgggaacca qqqaccccct atcttqqtac cqqtcattgg atgtatcccc agctcatgcc 300
tgtgtctgtc ttggcccgtg tggtcaccct gtgttcatct ctctcccagc catggcctct 360
caaactgggg ttttcgtctc cctatqaqqq gqtcctqqta tgtacgcgtt cggtgggccc 420
geggtgeatg teteceggtg cagtgeatge tggggtteee tggggeeetg ggeeeetegt 480
aggatagaca gagcotqtee taacetteeg qaaqtqcatg ctggggaggc cccttgcctg 540
ctgaccttct gtgctcagga cqactaatcg qccacatgac caccactctg tcccatggga 600
                                                                  637
ttcctagaga agtctcacta agagcccagc acactca
<210> 3
<211> 2693
<212> DNA
<213> Homo sapiens
<220>
<221> unsure
<222> (2266)..(2512)
<220>
<221> unsure
<222> (586)
```

```
<220>
<221> unsure
<222> (1480)
<220>
<221> unsure
<222> (1532)
<220>
<221> unsure
<222> (1562)..(1566)
<220>
<221> unsure
<222> (1569)
<220>
<221> unsure
<222> (1571)
<220>
<221> unsure
<222> (1631)
<400> 3
qctcctacaq ccqcatctqc qttaacataq catccctatq gccactqtct cccttgatcc 60
ccacagccat cctaggagaa aggcagaatg tcataatttg ctaaaaggga tgctgaggct 120
ctgggaggga aagggacttg cctaaagccc cagggtgaag cagcatctct ggactcccag 180
tocagtgate tigeceaata ettigetget tgeetatace eetetaaett ggteaacage 240
acatcacagg gcaagcccaa tecetgette attittatat atgggegetg gtecacagee 300,
ccactctcca qccatttqqa aacaaaaaca qatgctattg ttcttcctta gagaacgtgg 360
ccagtggaga cggcacactg gaaatcagag tgaatgttct tgaaagaggg tcacgggtca 420
acaaggccca gccaaaggat gcagtagaac cattttcctt agaaatcttt gggagtgaag 480
taggetteag ceactaceea tecetgeeet tgeggetace actaceeeat tagtttagae 540
agggtcgggc ggggaggggt gtggagaaga aatgagcttg cctgtngccc ccaggctccc 600
tetgteetag etcaggtetg ggtgeeatte tttacacteg tgtgeteget cacgeacaca 660
tcacacacct tgctggtcac acagtcacag actcgcctct gctcctgtgg tccagtggcc 720
ggacaccccc tgggatggct caaaggaqtc aggacttgga agtggggaca tcagggtagc 780
tgaaggaaat ccacacacce agagcatete ggagtteaga eteteagace tgaagtagge 840
gcccccggga ctqqqctaqq aqttqqacqq aatqqaggat ggaggacagc gagaagaaag 900
gaagagaaat gcaaagtgtg ggcagccgcc aagagtgaaa atagagggaa gtgtcatgca 960
agtgctggac agaaggcggc aggtgggacg agcccacag cccctcctc aaaaacgacc 1020
acctccagga ctcagtgatc cctggggggc aggctctgcc agccctcggc cacacgtggc 1080
tccggcaccc atggtcccag tgccttggat ggagacggcc agttctggcg gccagatgtg 1140
gtgctctgga atccagtccc atttccttcc tggccacgcc tgttccagcg gcctctttgg 1200
ctqcattcaq cccctactta cctqqqqacc ccqqctgggq cacaagagca ccaggggggt 1260
agggcccaaa gggatcaggg gaagcctctg gcctggaggg tatggggcac gcttccccaa 1320
```

<222> (52)

```
gggcggaccc qqcaqqaqqa aqcccaqqaq ctggqtcctg ccgcccaqqa gctgggccct 1380
gccacccagg ccgggctagg gacatggcag ggcctgggca tcctgacgct ggacttgggc 1440
gacctgggag gcacaqqqaq qqqaqaqatq ggcggcccn acccagcgca gtgccggcca 1500
caccccaagg cggttgccag agcttaaggc cnggccccag caggagaaca tcccagctcc 1560
annnnncene neegeageea gtgeteettg teaageteee eeegteaete eaggtgggag 1620
ccacccggt nagggggtgt gccacttgcc cccagggcac tcctctgggc atcccgggtg 1680
ggggattttg gggccgtggg gggcagtctc tggtacctgt gtgcgtcagg gatgctctqc 1740
acctgcaacc aggtgtcgtc cacgggeggg ggcatgggca tggtgacagt ggtcctgttg 1800
atgtcaccga tgatgctgag cgcctccttc agcgcgtggt gcatgtgcag catctcgtcg 1860
tgctgctgtg cctgctctgc caactcctcc atcagtgtgt tctggttccc acatgagtac 1920
atattggcca gcgctccga gatgatgaac tccggggtct gagagtgggc aaacagggaa 1980
gaaggttggg acctggtgcc tgtgccgccc tggctgcctt gctgggccct tctgggactg 2040
tgcgctggac ttggagccc ttggagtatg gcttttcaca cgggcttcta taccgcttcg 2100
actggaagat ccacctccc actgcctttt ctcactcaga tggggacacc gaggtccaga 2160
ggaaaagaca cctgtcaaat gtcacagatc tgggagggga cttaagacct atcatgccaa 2220
gaggacacct gtctactcag tttttttttg gtggggcggg gggcgnnnnn nnnnnnnnn 2280
agttgatgcc tggatacagg agctctgtgg gtgggagtga gacaaaacac agggtcctga 2580
gctctgggga ccaagcaatg tcctctggtg aaaaaaatcc tggacttgct ggcagaagat 2640
ttgcctctta cttgccatgt gctctgaata catttacctg ccctctggga aaa
                                                         2693
<210> 4
<211> 292
<212> DNA
<213> Homo sapiens
<220>
<221> unsure
<222> (284)
<400> 4
aagaatatga gatttgctta gaaatgaagg actggaagga gcccacagag tratttttta 60
aactatccag taaggcttag agggtttcaa tcagaaatat gtgttagggg aaaaaatgca 120
ctttttctat attaaaaaat attattttct tcttttaaat gtaaagcatt cctattgtga 180
agaattgaga aaatacagaa aagtacaaag aaaaacatta cctacaactc caccatccgt 240
gattatcact gttcacattt gtggctcatt tttcagtatk tctnttattt aa
<210> 5
<211> 2694
<212> DNA
<213> Homo sapiens
<220>
<221> unsure
```

```
<221> unsure
<222> (74)
<220>
<221> unsure
<222> (76)
<220>
<221> unsure
<222> (80)
<220>
<221> unsure
<222> (92)
<220>
<221> unsure
<222> (97)
<220>
<221> unsure
<222> (123)
<220>
<221> unsure
<222> (132)
<220>
<221> unsure
<222> (173)
<220>
<221> unsure
<222> (217)
<220>
<221> unsure
<222> (257)
<220>
<221> unsure
<222> (2539)
<400> 5
```

<220>

tactatattg ctcagcattt ctaagtattc tctaagtgct ctttatttat gntttaaaat 60 agctctctta cccngntgcg ncgactagaa gancttgntt taggaaacaa tgaaatatat 120 5

```
aanttgccag antcaattgg agccctctta catctaaaag atctctggtt ggntggaaat 180
caactgtcag aattacctca ggaaatagga aatctgnaga acctgctgtg tttagatgtc 240
totgaaaaca qqttqqnaaq acttootgaa gaaatcagtg gootgactto attaacggat 300
ttagtcattt cccagaactt attagaaacg attccggatg gcattggaaa actaaagaaa 360
ctgtcaatct tqaaggtqqa tcaqaataga ctcacacagt tgcctgaagc agttggggaa 420
tgtgaaagtc tcactgagtt agttcttaca gaaaatcagc tcctgaccct gcctaaaagc 480
attggaaaac taaagaagtt gagcaacttg aatgcagaca gaaataaatt agtgtcctta 540
ccaaaagaga teggegggtg etgeageete aetgtgttet gtgtaegtga caacagaeta 600
actoggatac ctgcagaggt gtcacaggca acagaacttc atgtcctgga tgtggcaggg 660
aacaggttgc tgcatctacc tttatccctg actgccttga agttgaaggc tctgtggcta 720
totgacaaco agtoccagoo cotgottaca ttocagacag acacagacta caccacagga 780
gagaagattt taacctgtgt cttacttcct cagctgcctt ctgaacctac ttgtcaagag 840
aatctgcctc gctgtggtgc actggagaac ttggtaaatg atgtctctga tgaagcctgq 900
aacqaqcqtq ctqtcaacaq aqtcaqtqcq atccqatttq tggaggatga gaaagatgaa 960
qaaqacaatq aqacqaqaac acttctaaqq cqaqccactc cacacccagg ggagttaaag 1020
cacatgaaaa agacagtgga gaatttacgg aatgacatga atgctgctaa aggactggac 1080
tcaaacaaaa acgaggtcaa tcatgccatt gaccgagtga ccacttctgt gtagagtttc 1140
acctccaagt tttacctcct gtgtcttcct ctgctgtcga gacgttcctg tctgcttccc 1200
gggagcctca cgtgctcctt gtcctaacca gcccccgcgc gccatcttcc cgtggagtgt 1260
ggggaagctg ctgtctccca ggaagtgcct tactcatccc gcaaccagtc agcgcaccag 1320
tggtctcccq qtgtgatttt ttttttttt aatttcagtt gtttgtaata agtagaatac 1380
actactgtaa acatacgacc tttgtttttg tcttatgttg gggtaaagga aagcaggaag 1440
qqqaattttt atcctcctcc cttccqtaaa qtqctqqqat attttqaatc ccccaagttc 1500
ccttggacct actgatgaga gatagtttta tgtatgggga aaaatggata ctttttaaac 1560
cttttttggc agctcagatg gtgtaaattt taaaattttg tataggtatt tcataacaaa 1620
aatatgtatt tottttttgt tattttatot tgaaaacggt acatatttta gtatttgtgc 1680
agaaaaacaa gtcctaaagt atttgttttt atttgtacca tccacttgtg ccttactgta 1740
tcctgtgtca tgtccaatca gttgtaaaca atggcatctt tgaacagtgt gatgagaata 1800
ggaatgtggt gttttaaagc agtgttgcat tttaatcagt aatctacctg gtggatttgt 1860
ttttaaccaa aaagatgaat tatcaatgat ttgtaattat atcggttgat tttttttgaa 1920
aagatgaacc aaaggatttq actgctaata ttttattcct tacacttttt ttctgaataa 1980
gtctctcata atgagtgcag tgtcagactg tgcctactct gatggtatgt gccatttgta 2040
aaataaaata gagcagaaaa acacaaaaag agaacactgg ttcagacatt cagtgggcaa 2100
gtaaattatg gactgcaaaa taatgatttt tattcaagaa agctttaaaa gttttatatc 2160
cagatataca accacaataa agcaaaataa cctactatca aaatagaaat qttqctatct 2220
ttataagtgc aatttaattt gtaaatagag tttgaatcaa agtatcacaa aatactgctt 2280
caaqatttaa ttttaaatct qctaatttaa qqqatattqq qaaaaqtttt qqtqtqtttc 2340
tgttgatttc ttttttgtat gctgtgataa aagagaaatg aaaagtgcca gtcactgtgt 2400
ggtgtctagg aaaatcatat atatttttt ctccaagaaa taaattcatc ctggacattg 2460
gccatacage tttttaaaat tattactttg tatgttcaag tgatagcagg tagccaaatt 2520
ctttgacagt gtgctctgnt ctgttaaata tctaaattac ccgtcagttg tgagtgacct 2580
cctgtgggac ttgcattcac atggggcaga gcccagaatt gcctttgact ctggctagta 2640
attttgggtt gtggctatct ggccaattgg actccttata aacccgtctt caac
```

<210> 6

<211> 1335

<212> DNA

<213> Homo sapiens

```
<220>
<221> unsure
<222> (17)
<400> 6
tcatatagta ggaaganaag cacctaggtt tgaggccagg gctggctgct gtcagaacct 60
aggecetece etgeettget ceacacetgg teaggggaga gaggggagga aagecaaggg 120
aaqqqaccta actqaaaaca aacaaqctgg gagaagcagg aatctgcgct cgggttccgc 180
agatgcagag gttgaggtgg ctgcgggact ggaagtcatc gggcagaggt ctcacagcag 240
ccaaggaacc tggggcccgc tecteceee tecaggccat gaggattetg cagttaatec 300
tgcttqctct qqcaacaqqq cttqtaqqqq qaqaqaccag gatcatcaag gggttcgagt 360
gcaagectea eteccagece tggcaggeag ecetgttega gaagaegegg etaetetgtg 420
gggcgacget categeeece agatggetee tgacageage ceaetgeete aageegtgge 480
cgctacatag ttcacctggg gcagcacaac ctccagaagg aggagggctg tgagcagacc 540
eggacageca etgagteett eecceacece ggetteaaca acageeteec caacaaagac 600
caccgcaatg acatcatgct ggtgaagatg gcatcgccag tctccatcac ctgggctgtg 660
cgacccctca ccctctcctc acgctgtgtc actgctggca ccagctgcct catttccggc 720
tggggcagca cgtccagccc ccagttacgc ctgcctcaca ccttgcgatg cgccaacatc 780
accatcattg agcaccagaa gtgtgagaac gcctaccccg gcaacatcac agacaccatg 840
gtgtgtgcca gcgtgcagga agggggcaag gactcctgcc agggtgactc cgggggccct 900
ctggtctgta accagtctct tcaaggcatt atctcctggg gccaggatcc gtgtgcgatc 960
accegaaage etggtgteta cacgaaagte tgcaaatatg tggactggat ccaggagacg 1020
atgaagaaca attagactgg acccacccac cacageccat cacectecat ttecacttgg 1080
tgtttggttc ctgttcactc tgttaataag aaaccctaag ccaagaccct ctacgaacat 1140
tetttgggcc tectggacta caggagatge tgtcacttaa taatcaacct ggggttegaa 1200
atcagtgaga cctggattca aattctgcct tgaaatattg tgactctggg aatgacaaca 1260
cetggtttgt tetetgttgt atecceagee ecaaagacag etectgeeat atateaagtt 1320
                                                                   1335
tcaataaata tttct
<210> 7
<211> 1079
<212> DNA
<213> Homo sapiens
<220>
<221> unsure
<222> (268)
<220>
<221> unsure
<222> (688)
<220>
<221> unsure
<222> (700)
<400> 7
```

tttttgaaga	atgccctgca	aggcatcaac	tggaatgtgt	ttattaccaa	acaagacaga	60
agagaaccag	ggcctgactt	ggcagtggcc	ccaggctgca	tgggctcagg	taggctcaga	120
ccggccccag	gagtgggaga	gcccagagaa	gagggaaaaa	gagtagtggc	caggaggggt	180
ctggctggga	catgccactc	tgggccatca	gcttctggat	ccactcaaag	tggtggctga	240
tattggtgta	gacaccgggc	cgattggncc	gaccacagcc	cactccccag	ctcacgactc	300
caatctgata	ccacagtcca	ttcttgttac	aggccaaggg	tccacctgag	tcaccgaagc	360
aggcatcctt	cccgccttgg	gcattgccag	cacaaaccat	gtctccaaag	atgtccttgc	420
ggaaactgta	cttgaggaag	aggtggttgc	acatagagtt	gtttatgatg	gcgacctgaa	480
cttcctggag	ggtgtgggga	gatggcagtg	cctcatcctc	tttgatgtac	ccccagccag	540
tcacccagca	gtctgtccgg	ttctcaaact	caaatgtgga	ggcctggaga	cagatgggct	600
ggatgtgttt	agtgtaggtg	acaggtgcag	acagetteae	caaggcaatg	tcatagggtg	660
aattccccag	ttagcgaggg	ctcagatnga	tattcgatan	gaagtaacgg	gtgtagtagg	720
cctgcaggct	ccagaaggat	ggcatggaag	tcagctggcc	aaactggacc	atccacccgg	780
agggatcact	aaggtcacta	taggtttcaa	agcagtgcgc	cgccgtgagt	gcccagcggt	840
ggctgagcag	gctcactccg	catacgtggg	aatcccacag	gcgcaggctc	ccctgccacg	900
gccaacgccc	gagttcggcg	tcctctccac	ccacgatgcg	cgacgtgatg	acccgtcggc	960
cgcatggtcc	tgataagggc	gccgcctcct	gcgactccgg	cttcctgagt	ccagcccgag	1020
ccagcagcag	cgccagcagc	agcgccccgc	gegegeceat	ggcctcctct	cccgcggtg	1079